

PCT / GB 05 / 00544

REC'D 10 MAR 2005

WFO

PCT

PA 1277375

**THE UNITED STATES OF AMERICA**

**TO ALL TO WHOM THESE PRESENTS SHALL COME:**

**UNITED STATES DEPARTMENT OF COMMERCE**

**United States Patent and Trademark Office**

**January 31, 2005**

**THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.**

**APPLICATION NUMBER: 60/546,343**

**FILING DATE: February 20, 2004**

**PRIORITY  
DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

**By Authority of the  
COMMISSIONER OF PATENTS AND TRADEMARKS**



*Trudie Wallace*  
**TRUDIE WALLACE**

**Certifying Officer**

# PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53 (c).

		DOCKET NUMBER	T1651PV
INVENTOR(S)			
Given Name (first and middle [if any])	Family Name or Surname	Residence (City and either State or Foreign Country)	
Ulf H. Anthony J. Karel M. J. Jeremy P. David R. Sarah E. Deborah C.	Dolling Davies Brands Scott Lieberman Brewer Hammond	Westfield, NJ Hoddesdon, Hertfordshire, UK Jersey City, NJ Hoddesdon, Hertfordshire, UK Hoddesdon, Hertfordshire, UK Hoddesdon, Hertfordshire, UK Hoddesdon, Hertfordshire, UK	
<input type="checkbox"/> Additional inventors are being named on the separately numbered sheets attached hereto			
TITLE OF THE INVENTION (500 characters max)			
CHEMICAL PROCESS			
CORRESPONDENCE ADDRESS			
Direct all Correspondence to: Merck & Co., Inc. Patent Department - RY60-30 P.O. Box 2000 Rahway			
		<input checked="" type="checkbox"/> Customer Number	000210
STATE	New Jersey	ZIP CODE	07065
		COUNTRY	U.S.A.
ENCLOSED APPLICATION PARTS (check all that apply)			
<input checked="" type="checkbox"/> Specification	Number of Pages	14	<input type="checkbox"/> CD(s), Number
<input type="checkbox"/> Drawing(s)	Number of Sheets		<input type="checkbox"/> Other (specify)
<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76			
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT (check one)			
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees		FILING FEE AMOUNT (\$)	\$160.00
<input checked="" type="checkbox"/> The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: 13-2755			

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No.

☐ Yes, the name of the U.S. Government agency and the Government contract number are: \_\_\_\_\_

Respectfully submitted,

SIGNATURE John C. Todaro  
TYPED or PRINTED NAME John C. Todaro  
TELEPHONE 762-594-0125

Date 02/20/2004

REGISTRATION NO. 36,036  
(if appropriate)

NOTE: Mail to Mail Stop Provisional Application

EXPRESS MAIL CERTIFICATE	
DATE OF DEPOSIT	<u>February 20, 2004</u>
EXPRESS MAIL NO.	<u>EL989593355US</u>
I HEREBY CERTIFY THAT THIS CORRESPONDENCE IS BEING DEPOSITED WITH THE UNITED STATES POSTAL SERVICE AS EXPRESS MAIL "POST OFFICE TO ADDRESSEE" ON THE ABOVE DATE IN AN ENVELOPE ADDRESSED TO COMMISSIONER FOR PATENTS, P.O. BOX 1450, ALEXANDRIA, VA 22313-1450.	
MAILED BY	<u>John Schipani</u>
DATE	<u>2-20-04</u>

In Duplicate

TITLE OF THE INVENTION  
CHEMICAL PROCESS


This invention is in the field of synthetic organic chemistry, and in particular concerns a novel synthetic route to a particular class of 4,4-disubstituted cyclohexanepropanoic acids which have useful therapeutic properties, and which are intermediates in the synthesis of further compounds having therapeutic properties.

As disclosed in WO 03/018453, a particular class of cyclohexane derivatives have been found to have activity as inhibitors of the processing of amyloid precursor protein (APP) by  $\gamma$ -secretase into the  $\beta$ -amyloid peptide. Since the secretion of  $\beta$ -amyloid is believed to play a primary role in the onset and progression of Alzheimer's disease, the said cyclohexane derivatives are useful in the treatment and/or prevention of Alzheimer's disease.

Included within the aforementioned class of cyclohexane derivatives are cyclohexanepropanoic acids, and salts, esters and amides derived therefrom, in which the carbon atom in the 4-position of the cyclohexyl ring is bonded to an aryl or heteroaryl group and also to an arylsulfonyl or heteroarylsulfonyl group. Said cyclohexanepropanoic acids exist in two stereoisomeric forms in which the propanoic acid group and the aryl- or heteroarylsulfonyl group are *cis* or *trans* with respect to the cyclohexane ring. Only the *cis* isomer has the requisite biological activity. The synthetic routes disclosed in WO 03/018453 and in US2003/0114496 involve condensation of a suitable 4,4-disubstituted cyclohexanone with an ylide and reduction of the resulting olefin. Isolation of the desired *cis* isomer requires the use of non-scaleable separation methods, or of expensive asymmetric reducing reagents and energy-intensive cryogenic conditions which render factory-scale preparations uneconomic.

There is therefore a need for an efficient, stereoselective synthesis of said cyclohexanepropanoic acids, amenable to execution on a large scale.

According to the invention there is provided a process for the preparation of a compound of formula I:





(1)

wherein R represents H or an alkali metal, Ar<sup>1</sup> represents 4-chlorophenyl and Ar<sup>2</sup> represents 2,5-difluorophenyl;

comprising the steps of:

(a) stirring a mixture of a *cis*-sulfide of formula (2) and a *trans*-sulfide of formula (3):



5 with 4-chlorobenzenethiol in an acidic medium in which said mixture of sulfides is partially soluble, causing preferential crystallisation of *cis*-sulfide of formula (2);

(b) collecting the *cis*-sulfide of formula (2);


(c) oxidising the *cis*-sulfide of formula (2) to the corresponding sulfone; and optionally

(d) neutralising the product of step (c) with alkali.

10 Suitable identities for R include H, Na, K and Li, but R is preferably H or Na.

The key step in the process defined above is the isolation of a sulfide of formula (2) in which the sulfide group and the propanoic acid group are in the requisite *cis* configuration. Thereafter, the sulfide group may be oxidised to the required sulfone by conventional means, and (if desired) the propanoic acid functionality converted to the corresponding alkali metal salt such as the sodium salt by neutralisation with the appropriate alkali, or to a suitable ester or amide derivative as described in WO 03/018453.

15 It is surprisingly found that the *cis*-sulfide of formula (2) is significantly less soluble than its *trans* isomer (3) in a wide variety of solvents, and hence can crystallise preferentially from a mixture of the two isomers. Furthermore, the isomers are believed to interconvert under acidic conditions via an olefin of formula (4):



Hence, in accordance with step (a) of the above-defined process, when a mixture of sulfides (2) and (3) is stirred in an acidic medium in the presence of excess Ar<sup>1</sup>SH (i.e. 4-

chlorobenzenethiol), essentially quantitative conversion of (3) to (2) may be achieved, with crystallisation of the latter in a high state of purity.

The nature and volume of the acidic medium must be such as to enable appreciable dissolution of the initial mixture of sulfides, otherwise the interconversion of the isomers becomes impractically slow (or non-existent). Generally speaking, increasing solubility of the sulfide mixture in the acidic medium leads to a higher rate of interconversion, but this may be at the expense of a reduced yield of crystalline product, with significant amounts of product remaining in solution. An initial solubility of about 0.1 – 1.0%w/v has been found to provide a suitable compromise, but values outside this range may be used if desired.

Addition of excess 4-chlorobenzenethiol to the system reduces the equilibrium concentration of the olefin (4) and hence increases the yield of sulfide (2). In at least some solvents, the presence of excess 4-chlorobenzenethiol also increases the solubility of the sulfides (2) and (3) and/or the solubility differential between the sulfides (2) and (3). The amount of excess 4-chlorobenzenethiol is not critical, but about 2 – 3 molar equivalents are typically used.

In one embodiment the acidic medium comprises a solvent and an acid. In principle, any solvent may be used in which a solubility differential exists in favour of the trans-sulfide (3). Examples include n-heptane, methylcyclohexane, trifluoroethanol, hexafluorobenzene, trifluorotoluene, hexafluoropropan-2-ol, acetonitrile and mixtures thereof. Others may be identified by simple experimentation. Particularly preferred solvents include hexafluoropropan-2-ol in admixture with another perfluorinated solvent such as perfluorohexane or perfluorinated 2-butyltetrahydrofuran (FC-75), e.g. a 1:1 v/v mixture of hexafluoropropan-2-ol and FC-75.

The acid is typically a strong organic acid such as a sulfonic acid or a perfluorinated carboxylic acid. Preferred acids include trifluoroacetic acid, benzenesulfonic acid, camphorsulfonic acid and C<sub>1-4</sub>alkylsulfonic acids in which one or more of the carbon atoms may optionally be perfluorinated. Particularly preferred acids include trifluoroacetic acid (TFA), trifluoromethanesulfonic acid (triflic acid) and methanesulfonic acid (MSA). The quantity of acid present is typically about 10 mole% (based on the sulfide mixture) in the case of triflic acid, and typically about 25 mol% in the case of MSA. Larger quantities may be used if desired, but no particular advantage ensues.

In a second embodiment, the acidic medium consists primarily of the acid itself, if necessary diluted with water to adjust the solubility of the sulfides to a suitable level. A preferred acid for use in this embodiment is MSA, e.g. containing from about 5 to about 15%v/v water, preferably about 9 to about 11%v/v water.

The mixture of sulfides (2) and (3) is preferably stirred in the acidic medium for at least one hour, more preferably for at least 10 hours, such as overnight. Said stirring may be


conducted at ambient or elevated temperatures. When hexafluoropropan-2-ol in admixture with another perfluorinated solvent is used as the acidic medium, overnight stirring at ambient temperature is suitable. When aqueous MSA solvent is used as the acidic medium, overnight stirring at about 35 to 45°C is suitable.

5 After cooling (if necessary) the *cis*-sulfide of formula (2) is collected, e.g. by filtration. If desired, the filtrate may also be collected and recycled. This may be important when the acidic medium comprises one or more expensive fluorinated solvents. As an alternative to filtration, the *cis*-sulfide (2) may be collected by extraction into a solvent which is immiscible with the acidic medium. For example, when the acidic medium is aqueous MSA, extraction into ethyl  
10 acetate or isopropyl acetate may be used, the sulfide being isolated by subsequent evaporation of the extract to dryness. Prior to collecting the *cis*-sulfide, the mixture may be diluted with an antisolvent for the sulfide to promote further crystallisation of same. For example, when MSA containing 5-15%v/v water is used as the acidic medium, it is advantageous to dilute the medium with an approximately equal volume of water prior to collection of the *cis*-sulfide. The material  
15 thus obtained is typically more than 90% pure. If desired, further purification may be effected by recrystallisation (e.g. from acetonitrile or another solvent system) to provide *cis*-sulfide (2) that is essentially free from the *trans* isomer (3).

Step (c) of the process according to the invention, namely oxidation of the *cis*-sulfide (2) to the corresponding sulfone, may be effected by any of the oxidising agents conventionally used  
20 for the conversion of sulfides to sulfones, such as hydrogen peroxide, peroxyacetic acid or *m*-chloroperoxybenzoic acid (see, for example, Trost and Fleming, *Comprehensive Organic Synthesis*, 1993, Vol 7, 766). A preferred procedure comprises addition of about 3 equivalents of hydrogen peroxide to a solution of the sulfide in acetic acid, followed by heating at about 55-60°C for about 5 hours, cooling to ambient temperature and dilution with water to precipitate the  
25 sulfone product.


Optional step (d) of the process according to the invention, namely neutralisation of the carboxylic acid group, may be carried out according to published procedures (e.g. US2003/0114496). A suitable process involves addition of 2M aqueous sodium hydroxide to a solution of the acid in propan-2-ol, followed by distillation of the solvent.

30 The mixture of *cis*- and *trans*-sulfides used in step (a) of the process of the invention may be generated by any suitable means. In one embodiment of the invention, said mixture is generated by the reaction of 4-chlorobenzenethiol with an olefin of formula (4):



wherein  $\text{Ar}^2$  represents 2,5-difluorophenyl. Advantageously, said reaction is carried out in the acidic medium subsequently used in step (a) of the process of the invention, so that the mixture of *cis*- and *trans*-sulfides is generated *in situ*.

In another embodiment of the invention, said mixture is generated by the reaction of 4-chlorobenzenethiol with a carbinol of formula (5):



wherein  $\text{Ar}^2$  represents 2,5-difluorophenyl, said reaction being carried out in the presence of a Lewis acid, and the mixture of sulfides being isolated prior to carrying out step (a) of the process of the invention. In principle, any of the known Lewis acids may be used (e.g. those disclosed in "Lewis Acids in Organic Synthesis", ed. Yamamoto, pub. Wiley, 2000). Preferred examples include scandium triflate, boron trifluoride diethyl etherate, ethyl aluminium dichloride, zinc iodide, aluminium trichloride, tin tetrachloride, indium(II) chloride, indium(III) chloride, titanium tetrachloride, methyl aluminium dichloride and zinc triflate, of which boron trifluoride diethyl etherate is particularly preferred. The reaction may be carried out in an aprotic solvent such as dichloromethane or acetonitrile at reduced temperature (e.g. about -10 to -15°C).

As will be readily apparent to those skilled in the art, carbinols represented by formula (5) exist as *cis* and *trans* isomers. Structural formula (5) encompasses both of said isomers, either as pure compounds or as mixtures in any proportion. Both isomers are equally useful in the invention, and carbinols (5) are typically generated and used as an approximately 1:1 mixture of *cis* and *trans* isomers.

In a preferred embodiment of the invention, said mixture of *cis*- and *trans*-sulfides is generated by the reaction of 4-chlorobenzenethiol with a carbinol mixture of formula (5) in the acidic medium subsequently used for step (a) of the process of the invention.


In a typical procedure in accordance with the invention, 4-chlorobenzenethiol is charged to a suitable reaction vessel and methanesulfonic acid added in sufficient quantity to dissolve most of the thiol when heated to about 40°C (e.g. about 2.4 litres of acid per mole of thiol). Water (from about 8% to about 12% v/v with respect to the acid) is then added slowly with stirring and cooling so as to maintain a temperature of about 30-40°C. Thereafter, the carbinol (5) and/or olefin (4) (about 0.8 – 0.9 equivalents relative to the thiol) is added to the mixture. This results in rapid consumption of the carbinol and formation of a mixture comprising olefin (4) and *cis*-sulfide (2) together with its *trans* isomer. Aging of the above-described reaction mixture results in a selective accumulation of the desired *cis*-isomer, such that dilution with

water, filtration and washing provides a high yield of crude product comprising >90% pure *cis*-sulfide (2). Reproducibility is facilitated by the addition of small amounts of pure *cis*-sulfide (2) as a seed prior to the aging step. Using this protocol, product of >99% purity is routinely obtainable via a single recrystallisation of the crude product. The quantity of seeding material is not critical, about 1% by weight relative to the carbinol (5) and/or olefin (4) being sufficient. Aging is carried out for at least 1 hour, preferably at least 3 hours and most preferably at least 10 hours (e.g. overnight). Aging can be conducted at ambient temperature but is preferably conducted by stirring at moderately elevated temperatures, for example in the range of about 35°C to about 45°C.

At the end of the aging period, the mixture is cooled to ambient temperature and the crude product isolated, e.g. by dilution with water, filtration and washing of the crude solid product with further water. Alternatively, the diluted aqueous mixture can be extracted with a water immiscible organic solvent (e.g. ethyl acetate or isopropyl acetate) which is subsequently evaporated to dryness to provide the same crude product. The crude *cis*-sulfide (2) is preferably further purified by recrystallisation from a suitable solvent such as acetonitrile.

In further embodiments of the invention, the carbinol mixture of formula (5) is prepared by:

(a) conversion of carboxylic acid (6a) to magnesium salt (6b):



(6) (a)  $R' = H$   
(b)  $R' = MgX$

(b) reaction of (6b) with  $Ar^2-M'$ ; and

(c) treatment of the resulting product with acid;

wherein  $M'$  represents Li,  $MgX$  or  $CeX_2$ ;  $X$  represents Cl, Br or I; and  $Ar^2$  represents 2,5-difluorophenyl.

$X$  preferably represents Cl.  $M'$  preferably represents Li.

Before reaction with  $Ar^2-M'$ , the carboxylic acid (6a) is converted to the magnesium salt (6b), e.g. by treatment with one equivalent of  $R^2MgX$ , where  $R^2$  represents  $C_{1-6}$ alkyl, preferably  $C_{1-4}$ alkyl such as methyl, ethyl, n-propyl or isopropyl (especially isopropyl), and  $X$  represents Cl, Br or I, preferably Cl. This process may be carried out in an ether solvent (preferably THF) at reduced temperature, e.g. below -60°C, under nitrogen or other inert gas. Alternatively, (6a) may be treated with solid  $MgX_2$  in the presence of a base such as triethylamine, e.g. in THF.



The reagent  $\text{Ar}^2\text{-M}'$  may be prepared by conventional means. For example, treatment of  $\text{Ar}^2\text{-H}$  with a slight excess of n-butyllithium in the presence of one equivalent of N,N,N,N-tetramethylethylenediamine in THF below  $-60^\circ\text{C}$  under nitrogen provides a solution of  $\text{Ar}^2\text{-Li}$  suitable for use in the process (cf. Ong et al, *J. Med. Chem.* **1981**, 24, 74 and Bridges et al, *J. Org. Chem.* **1990**, 55, 773). Similarly, treatment of  $\text{Ar}^2\text{-Br}$  with one equivalent of  $\text{R}^2\text{MgX}$  in THF below  $-10^\circ\text{C}$  under nitrogen provides a solution of  $\text{Ar}^2\text{-MgX}$  suitable for use in the process, where  $\text{R}^2$  and  $\text{X}$  have the same definitions and preferred identities as before (cf. Abarbri et al, *Tetrahedron Lett.* **1999**, 40, 7449). Addition of a THF solution of  $\text{Ar}^2\text{-MgX}$  to a slurry of  $\text{CeX}_3$  in THF at about  $-10\text{--}0^\circ\text{C}$  provides a solution of  $\text{Ar}^2\text{-CeX}_2$  suitable for use in the process.

Reaction of (6b) with  $\text{Ar}^2\text{-M}'$  is typically carried out in THF at about  $-60$  to  $-20^\circ\text{C}$  (when  $\text{M}' = \text{Li}$ ) or about  $-30$  to  $-20^\circ\text{C}$  (when  $\text{M}' = \text{MgX}$ ) or about  $0^\circ\text{C}$  when  $\text{M}' = \text{CeX}_2$ . Conventional work-up (e.g. quenching with acetic acid at low temperature, warming to ambient, dilution with dilute hydrochloric acid and extraction into toluene) provides the crude carbinol mixture of formula (5) which may be further purified by recrystallisation (e.g. from a mixture of toluene and heptane).

The compound (6a) is known in the literature (Adkins et al, *J. Am. Chem. Soc.*, **1938**, 60, 1467; and Adkins et al, *J. Am. Chem. Soc.*, **1940**, 62, 2422). It is conveniently prepared by hydrogenation of 3-(4-hydroxyphenyl)propanoic acid or 3-(4-hydroxyphenyl)propenoic acid over  $\text{Rh}/\text{Al}_2\text{O}_3$  in isopropyl acetate at about  $70^\circ\text{C}$ , followed by oxidation of the resulting cyclohexanol with  $\text{NaOCl}$  in the presence of  $\text{RuCl}_3$  and acetic acid in isopropyl acetate. Details of these, and all other steps of the inventive process, are provided in the Examples section contained herein.

The compounds of formula (2), formula (4) and formula (5) are themselves novel, and individually constitute further aspects of the invention. The compounds of formula (4) are formed in situ when compounds of formula (5) are mixed with aqueous methanesulfonic acid and  $\text{Ar}^1\text{-SH}$ . However, they may be isolated in pure form if desired by treatment of carbinols (5) with acid (e.g. trifluoroacetic acid) in the absence of  $\text{Ar}^1\text{-SH}$  in an aprotic solvent such as dichloromethane.

The products of the novel process disclosed herein have an activity as inhibitors of the processing of APP by  $\gamma$  secretase, and are therefore useful in the treatment or prevention of disorders involving excessive secretion and/or deposition of  $\beta$ -amyloid, in particular Alzheimer's disease.

The products formed via the inventive process may be used to prepare pharmaceutical compositions comprising one or more of the said products or pharmaceutically acceptable salts thereof and a pharmaceutically acceptable carrier. Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or

suspensions, metered aerosol or liquid sprays, drops, ampoules, transdermal patches, auto-injector devices or suppositories; for oral, parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, such as the conventional tableting ingredients known to those skilled in the art, e.g. as described in WO 01/70677, and formed into unit dosage forms. Typical unit dosage forms contain from 1 to 100 mg, for example 1, 2, 5, 10, 25, 50 or 100 mg, of the active ingredient. Tablets or pills of the composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action, as described, for example, in WO 01/70677.

The liquid forms in which the compositions may be incorporated for administration orally or by injection include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils, as described in WO 01/70677.

For treating or preventing Alzheimer's disease, a suitable dosage level of the active ingredient is about 0.01 to 250 mg/kg per day, preferably about 0.05 to 100 mg/kg per day, and especially about 0.1 to 50 mg/kg of body weight per day. The compounds may be administered on a regimen of 1 to 4 times per day. In some cases, however, dosage outside these limits may be used.

Assays for determining the level of activity of the relevant compounds towards  $\gamma$ -secretase are disclosed in WO 01/70677 and in *Biochemistry*, **2000**, 39(30), 8698-8704. See also, *J. Neuroscience Methods*, **2000**, 102, 61-68.

## EXAMPLES

### Example 1

#### 3-(4-oxo-cyclohexyl)propanoic acid

A mixture of 3-(4-hydroxyphenyl)propenoic acid (3.33Kg), 5% Rh/Al<sub>2</sub>O<sub>3</sub> catalyst (248g) and isopropyl acetate (IPAc) (30L) was hydrogenated (200 psi) at 70°C for 12 hours, then cooled and filtered.

To the resulting solution of 3-(4-hydroxycyclohexyl)propanoic acid was added RuCl<sub>3</sub> (42g) and acetic acid (5.6Kg). After cooling to 5°C, aqueous NaOCl (10-13%, 26Kg) was added slowly with stirring, maintaining the temperature at 5°C. After stirring a further 3 hours at 5°C, 3N HCl (20L) and IPAc (15L) were added, the layers were separated and the aqueous layer extracted with IPAc (2x20L). The combined organics were washed with 2M aqueous NaHSO<sub>3</sub> (10L), dried (MgSO<sub>4</sub>), treated with charcoal and filtered. The resulting solution was concentrated, and the solvent switched to heptane:IPAc 2:1 (20L). This solution was seeded and the solvent switched

to 9:1 heptane:IPAc, giving the product as a white solid which was collected, washed with heptane and dried under nitrogen.

### Example 2

5 *cis* and *trans*-3-[4-(2,5-difluorophenyl)-4-hydroxycyclohexyl]propanoic acid  
(Method 1)

*N,N,N,N*-tetramethylethylenediamine (8.8 ml, 57.8 mmol) was diluted with THF (70 mL) and purged with N<sub>2</sub> (3x) before cooling to -70°C. <sup>n</sup>Butyllithium (25.2 mL, 57.8 mmol) was then added dropwise maintaining the temperature below -60°C. After aging 10 minutes at -65°C,  
10 1,4-difluorobenzene (6 g, 5.4 mL, 52.6 mmol) was added dropwise, maintaining the temperature below -60°C, and the resulting solution was aged at -60°C for 90 minutes.  
In a separate vessel, 3-(4-oxo-cyclohexyl)propanoic acid (5.38 g, 31.6 mmol) in THF (80 mL) was purged with N<sub>2</sub> (3x) before cooling to -65°C. Isopropylmagnesium chloride (2M in THF) (15.5 mL, 31.0 mmol) was added dropwise over 30 min keeping the temperature below -60°C.  
15 The resultant solution was warmed to -10°C, then added slowly to the aryllithium solution prepared above, maintaining the temperature below -60°C. Upon complete addition, the temperature was allowed to warm to -20°C slowly. The thick slurry was quenched by the dropwise addition of acetic acid (6 mL, 104 mmol) below 0°C and allowed to warm to ambient. 2M HCl (140 mL) was charged to the reaction mixture followed by toluene (140 mL). The  
20 layers were cut and the organic layer was washed with H<sub>2</sub>O (1 x 100 mL).  
Evaporation in vacuo to dryness afforded crude product. This was redissolved in toluene (45 mL) at 80°C then allowed to cool to 45°C and seeded, at which point a seedbed formed.  
Heptane (45 mL) was added dropwise over 30 min at 45°C then the slurry was allowed to cool to ambient. After 19 hours, the slurry was filtered, washed with 2:1 heptane:toluene (15ml) then  
25 heptane (15ml) and dried under vacuum at 50°C.

### Method 2

1-Bromo-2,5-difluorobenzene (157.3 g, 0.815 mol) was diluted with THF (500 mL) and purged with N<sub>2</sub> (3x) before cooling to -30°C. Isopropylmagnesium chloride (2.0M in THF) (400 mL,  
30 0.80 mol) was then added dropwise maintaining a temperature below -10°C. The resultant opaque solution of 2,5-difluorophenylmagnesium grignard was aged 1.0 hr at -10°C.  
In a separate vessel, 3-(4-oxo-cyclohexyl)-propionic acid (85.1 g, 0.50 mol) in THF (1000 mL) was purged with N<sub>2</sub> (3x) before cooling to -58°C. Isopropylmagnesium chloride (2.0M in THF) (240 mL, 0.48 mol) was added dropwise maintaining a temperature below -55°C. The resultant  
35 solution was aged for 0.5 h at -55 to -50°C then warmed to -30°C over 15 min and held at this

temperature a further 0.5 h. This solution was added slowly to the solution of difluorophenylmagnesium grignard cooled to  $-30^{\circ}\text{C}$ , so that the temperature remained in the range  $-30$  to  $-20^{\circ}\text{C}$ . The resultant thick slurry was aged for 0.75 h below  $-10^{\circ}\text{C}$ . The mixture was quenched with acetic acid (86 mL, 1.50 mol) below  $5^{\circ}\text{C}$ , warmed to  $10^{\circ}\text{C}$ , then treated with 1M HCl (750 mL) and toluene (750 mL), aged at  $25^{\circ}\text{C}$  for 10 min and then the layers cut. The organic layer was washed with  $\text{H}_2\text{O}$  (1 x 1250 mL) and then extracted with 1M NaOH (1250 mL). The NaOH layer was stirred with toluene (1500 mL) before 2M HCl (700 mL) was charged at  $22^{\circ}\text{C}$  and the mixture agitated for 5 min. The toluene layer was washed with  $\text{H}_2\text{O}$  (2 x 1000 mL) then filtered and evaporated in vacuo to dryness. The crude product was purified as described in method one.

### Example 3

#### cis- and trans-3-[4-(4-chlorophenyl)thio-4-(2,5-difluorophenyl)cyclohexyl]propanoic acid

4-Chlorobenzenethiol (15.67g, 108mmol) in dichloromethane (46ml) under nitrogen was cooled to  $-15^{\circ}\text{C}$  and treated with boron trifluoride diethyl etherate (14.3g, 101mmol). A solution of the carbinol mixture from Example 2 (22g, 77mmol) in dichloromethane (245ml) was added slowly, keeping the temperature in the range between  $-9$  and  $-15^{\circ}\text{C}$ . After stirring a further two hours, the solvent was evaporated, replaced with acetonitrile, and this also evaporated. The residue was slurried with acetonitrile (30ml) and water (100ml) added. The resulting solid was filtered, washed with water, dried at  $40^{\circ}\text{C}$  *in vacuo* and recrystallised from heptane. HPLC analysis indicated a 1:1 mixture of *cis* and *trans* isomers.

### Example 4

#### cis-3-[4-(4-chlorophenyl)thio-4-(2,5-difluorophenyl)cyclohexyl]propanoic acid

The product from Example 2 (100mg) and 4-chlorothiophenol (70mg) were added to a mixture of hexafluoropropan-2-ol (0.5ml) and FC-75 (0.5ml), followed by triflic acid (8 $\mu\text{l}$ ). The resulting slurry was stirred overnight at ambient temperature, then filtered. HPLC analysis of the solid product indicated a mixture of 98.5% of the *cis* isomer and 1.5% of the *trans* isomer had been obtained.

### Example 5

#### cis-3-[4-(4-chlorophenyl)thio-4-(2,5-difluorophenyl)cyclohexyl]propanoic acid

To a stirred mixture of 4-chlorothiophenol (65.0g, 0.449mol) and methanesulfonic acid (1.07L) under nitrogen was slowly added water (118ml), with cooling to maintain a temperature of  $30$ - $40^{\circ}\text{C}$ . The carbinol from Example 2 (106.8g, 0.374mol) was added in a single portion, then after

10 minutes 1g of purified *cis*-sulfide was added as seed. The mixture was stirred overnight at 36-42°C, cooled, and diluted slowly with water (1.07L) with continued cooling to maintain a temperature of 22-39°C. After aging a further 30 minutes, the white solid was collected by filtration, washed with water and dried at 40°C under vacuum. The crude product was slurried with acetonitrile (approximately 300ml per 100g of product), heated to reflux until dissolution was complete, then allowed to cool. The resulting crystalline material was collected, washed with further acetonitrile and dried 40°C under vacuum. Final purity was assayed as 98.7 wt%.

#### Example 6

##### *cis*-3-[4-(4-chlorophenyl)sulfonyl-4-(2,5-difluorophenyl)cyclohexyl]propanoic acid

The product from Example 5 (5.0g, 12.17 mmol) in acetic acid (50ml) was treated dropwise at 50°C with hydrogen peroxide (27.5 wt%, 4.07ml, 36.5mmol) and the mixture kept at 55-60°C for 4.75 hours. After cooling to ambient temperature, water (100ml) was added dropwise and the resulting slurry stirred a further 1 hour. The product was collected by filtration, washed with water and dried at 40°C *in vacuo*.

#### Example 7

##### *cis*-3-[4-(4-chlorophenyl)sulfonyl-4-(2,5-difluorophenyl)cyclohexyl]propanoic acid, sodium salt

Product from Example 6 (100g, 0.226mol) in isopropanol (2L) at 45°C was treated with 2M aqueous sodium hydroxide (112ml). The resulting solution was distilled at atmospheric pressure, removing 1L distillate. Fresh isopropanol (1L) was added and the distillation repeated. After adding further fresh isopropanol (1L) and repeating the distillation again, the mixture was allowed to cool to ambient and aged a further 1 hour. The product was collected by filtration, washed with isopropanol and dried at 40°C *in vacuo*.


#### Example 8

##### [4-(2,5-difluorophenyl)cyclohex-3-enyl]propanoic acid

Trifluoroacetic acid (6.8ml, 87.9mmol) was added to a solution of the product of Example 2 (2.5g, 8.79mmol) in dichloromethane (40ml). The resulting mixture was stirred at ambient temperature overnight and then partitioned between water (50ml) and IPAc (150ml). The layers were separated. The organic layer was washed with water (50ml) and brine (50ml), dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was recrystallised from IPAc, and dried at 40°C *in vacuo*.


## WHAT IS CLAIMED IS:

1. A process for the preparation of a compound of formula (1):




- 5 wherein R represents H or an alkali metal, Ar¹ represents 4-chlorophenyl and Ar² represents 2,5-difluorophenyl;  
comprising the steps of:

- (a) stirring a mixture of a *cis*-sulfide of formula (2) and a *trans*-sulfide of formula (3):



(2)



(3)

10 with 4-chlorobenzenethiol in an acidic medium in which said mixture of sulfides is partially soluble, causing preferential crystallisation of *cis*-sulfide of formula (2);

- (b) collecting the *cis*-sulfide of formula (2);

- (c) oxidising the *cis*-sulfide of formula (2) to the corresponding sulfone; and optionally

- 15 (d) neutralising the product of step (c) with alkali.


2. A process according to claim 1 wherein said acidic medium comprises an acid selected from trifluoroacetic acid and C<sub>1-4</sub>alkylsulfonic acids in which one or more of the carbon atoms may optionally be perfluorinated.

3. The process according to claim 2 wherein the acid is trifluoroacetic acid, trifluoromethanesulfonic acid or methanesulfonic acid.

4. A process according to claim 2 wherein said acidic medium additionally comprises a solvent selected from n-heptane, methylcyclohexane, trifluoroethanol, hexafluorobenzene, trifluorotoluene, hexafluoropropan-2-ol, acetonitrile and mixtures thereof.


5. A process according to claim 1 wherein the acidic medium is methanesulfonic acid containing from about 5 to about 15 % water by volume.

6. A process according to claim 1 wherein the mixture of *cis*-sulfide of formula (2) and *trans*-sulfide of formula (3) is generated by reaction of 4-chlorobenzenethiol with an olefin of formula (4):




wherein Ar² represents 2,5-difluorophenyl, said reaction being carried out in the acidic medium used in step (a) of the said process.

7. A process according to claim 1 wherein the mixture of *cis*-sulfide of formula (2) and *trans*-sulfide of formula (3) is generated by reaction of 4-chlorobenzenethiol with a carbinol of formula (5):



wherein Ar² represents 2,5-difluorophenyl, said reaction being carried out in the presence of a Lewis acid, and the mixture of sulfides being isolated prior to carrying out step (a) of the said process.

8. A process according to claim 1 wherein the mixture of *cis*-sulfide of formula (2) and *trans*-sulfide of formula (3) is generated by reaction of 4-chlorobenzenethiol with a carbinol of formula (5):



wherein Ar² represents 2,5-difluorophenyl, said reaction being carried out in the acidic medium used in step (a) of the said process.


9. A process according to claim 6 or claim 8 wherein the acidic medium comprises an acid and hexafluoropropan-2-ol together with a co-solvent selected from perfluorohexane and perfluorinated 2-butyltetrahydrofuran.

10. A process according to claim 9 wherein the acid is trifluoromethanesulfonic acid.

11. A process according to claim 6 or claim 8 wherein the acidic medium is methanesulfonic acid containing from about 5 to about 15 % water by volume.

12. A process according to claim 7 or claim 8 wherein the carbinol of formula (5) is prepared by:

(a) conversion of carboxylic acid (6a) to magnesium salt (6b):



(6) (a)  $R' = H$   
(b)  $R' = MgX$

(b) reaction of (6b) with  $Ar^2-M'$ ; and


(c) treatment of the resulting product with acid;

wherein  $M'$  represents Li,  $MgX$  or  $CeX_2$ ;

$X$  represents Cl, Br or I; and

$Ar^2$  represents 2,5-difluorophenyl.


13. The compound of formula (5):



(5)

where  $Ar^2$  is 2,5-difluorophenyl.


14. The compound of formula (4):



(4)

wherein  $Ar^2$  is 2,5-difluorophenyl.

15. The compound of formula (2):



(2)

where  $Ar^1$  is 4-chlorophenyl and  $Ar^2$  is 2,5-difluorophenyl.